

Dawn Mission
Mars Gravity Assist
Preliminary Report for GRaND, v1.1

Prepared by Tom Prettyman

3-Mar-2009

Overview

- GRaND acquired neutron and gamma ray counting data during Mars Gravity Assist
 - Data acquired up to S/C entry into safe-mode (occurred after MCA at an altitude of 3000 km)
- GRaND got close enough to Mars to see neutrons and gamma rays:
 - Mars closest approach (MCA) was about 550 km
 - About 16 min were spent below 1000 km during which time the S/C was nadir-pointing to within about 10°
- The space background environment was quiet, ideal for science data acquisition
- GRaND was nominal during the encounter
- Preliminary results with comparisons to the Odyssey NS are presented


Operations and Data Analysis

- Instrument settings
 - 35s measurement intervals below 16000 km
 - Timestamp regression was not observed
- Data Volume
 - 295 science data records
- Level 1a data delivered to DSC
- Higher order data products under development
- Instrument was powered on successfully following MGA yesterday (2-Mar-09)
 - Continued data acquisition followed by power-off in DC024
 - Anneal operations deferred
 - Lessons learned (Should HV sequence be separate from Power ON and parameter set?)

Mars Closest Approach


18-Feb-2008, $L_S=212^\circ$ Mid Autumn in the Northern Hemisphere

Thermal




Seasons on Mars: Northern Hemisphere, T. H. Prettyman (LANL) and T. N. Titus (USGS)


Epithermal



Seasons on Mars: Northern Hemisphere, T. H. Prettyman (LANL) and T. N. Titus (USGS)



Seasons on Mars: Southern Hemisphere, T. H. Prettyman (LANL) and T. N. Titus (USGS)




Seasons on Mars: Southern Hemisphere


Trajectory (below 1000 km)

The seasonal cap in the northern hemisphere was thin.
At high northern latitudes, the neutron spectrometer
was primarily sensitive to the water-rich regolith.

GRaND subsatellite
points and Field of View




Counting Data




Neutron Counting Data

Nadir-pointing Li-glass
CAT1, ${}^6\text{Li}(n,t)$ peak



Fast neutrons
CAT4, nadir+sides




The counting data were averaged using a 5-point sliding window

Analysis Method: Simple Approach





Procedure:

- Correct GRaND time series data for altitude and S/C orientation
 - Divide by the solid angle subtended by Mars at the S/C
 - Divide by the cosine of the S/C angle relative to nadir
- Compare the corrected GRaND counting rate for each measurement to the Odyssey counting rate averaged over GRaND's footprint

Simulated, footprint-averaged Odyssey data prepared prior to Dawn/MGA




Comparison Measured GRaND counting data to Odyssey



Detailed Analysis

- Neutrons
 - Compare modeled counting rates based on neutron leakage spectra to GRaND data
 - Accounts for neutron decay, ballistic trajectories and relative motion of the S/C (ram effect)
- Gamma rays
 - Compare O and Fe counting rates to mapped ODY/GRS data available in the PDS
- Analysis will include implications for science at Vesta and Ceres
- Results to be presented at EGU
- Manuscript on GRaND performance during cruise and MGA in preparation



Conclusions

- Mars Gravity Assist was a tremendous success from the standpoint of acquiring useful GRaND data
- GRaND measured gamma rays and neutrons originating from Mars
- GRaND neutron counting rates are strongly correlated with footprint-averaged Odyssey data
- The GRaND data will be useful for calibration and mission planning
- Results presented here should be regarded as preliminary - additional work on data reduction and analysis methods is needed in order to provide accurate comparisons to Odyssey